arXiv:1404.5470v1 [math.GR] 22 Apr 2014

The Möbius Function of the Suzuki Groups, with Applications to Enumeration

Martin Downs and Gareth A. Jones
School of Mathematics
University of Southampton
Southampton SO17 1BJ, U.K.
G.A.Jones@maths.soton.ac.uk

Abstract

We compute the Möbius function for the subgroup lattice of the simple Suzuki group $Sz(q)$; this is used to enumerate normal subgroups of certain Hecke groups with quotients isomorphic to $Sz(q)$.

MSC classification: 20G41(primary); 05A99, 11A25, 14H57, 20B20, 20H10 (secondary).

Key words: Suzuki group, Möbius function, subgroup lattice, Hecke group.

1 Introduction

Hall's theory of Möbius inversion in groups [8] allows one to enumerate various objects associated with a given finite group G. For example, using this method one can compute the number of normal subgroups N of the free group F_k of finite rank k with $F_k/N \cong G$, or equivalently the number of orbits of $\mathrm{Aut}\, G$ on generating k-tuples for G. Indeed, both of these numbers are equal to

$$d_k(G) = \frac{1}{|\mathrm{Aut}\, G|} \sum_{H \leq G} \mu_G(H)|H|^k \tag{1}$$

where μ_G is the Möbius function on the lattice of subgroups of G, defined recursively by

$$\sum_{K \geq H} \mu_G(K) = \delta_{H,G}. \tag{2}$$

Here $\delta_{H,G}$ is the Kronecker delta function, equal to 1 or 0 as $H = G$ or $H < G$.) As examples of this, $d_2(G)$ gives

- the number of isomorphism classes of orientably regular hypermaps with automorphism group G;
- the number of regular unbranched coverings of the sphere minus three points (or of the torus minus one point) with covering group G;
- the number of regular dessins (in Grothendieck's terminology [6]) with automorphism group G.

Equation (1) is a particular case of the general result that the number $n_\Gamma(G)$ of normal subgroups N of a finitely generated group Γ with quotient isomorphic to a given finite group G is given by

$$n_\Gamma(G) = \frac{|\mathrm{Epi}(\Gamma, G)|}{|\mathrm{Aut}\, G|} = \frac{1}{|\mathrm{Aut}\, G|} \sum_{H \le G} \mu_G(H) |\mathrm{Hom}(\Gamma, H)|, \tag{3}$$

which arises from applying Möbius inversion to the obvious equation

$$|\mathrm{Hom}(\Gamma, G)| = \sum_{H \le G} |\mathrm{Epi}(\Gamma, H)|. \tag{4}$$

A similar principle applies to the enumeration of torsion-free normal subgroups of Γ with quotient G: one simply counts the smooth homomorphisms and epimorphisms $\Gamma \to H$, those preserving the orders of torsion elements.

Implementing equation (3), as for some other similar results, for a specific group G depends on knowing the value of $\mu_G(H)$ for each subgroup $H \le G$. Finding these values can be a laborious task, but once it has been done, equation (3) can be applied to G in many different contexts, depending on the choice of Γ (see [4, 12], for example). The function μ_G has been determined for certain groups G, including the simple groups $L_2(p) = PSL_2(p)$ for primes $p \ge 5$, by Hall in [8]. Subsequently the first author extended this in [1] to the groups $L_2(q)$ and $PGL_2(q)$ for all prime powers q: see [2] for full details for $L_2(2^e)$ and a statement of results for $L_2(q)$ with q odd, and [4] for some applications.

The objective of this paper, which is based on an earlier preprint [3] by the first author, is to calculate the Möbius function μ_G for the family of simple groups $G = Sz(q) = {}^2B_2(q)$, where $q = 2^e$ for some odd $e > 1$. These groups were discovered in 1960 by Suzuki [14, 15]. We describe G and its subgroups H in Section 2, and give the values of $\mu_G(H)$ in Table 1 in Section 2.4. Specifically, this table gives the values of $\mu_G(H)$ and $|N_G(H)|$ for a set $\mathcal{T}$ of representatives H of the conjugacy classes of subgroups on which μ_G can take non-zero values. This information is sufficient for applications of equation (3): since $|\mathrm{Aut}\, G| = e|G|$, this now takes the form

$$n_\Gamma(G) = \frac{1}{e} \sum_{H \in \mathcal{T}} \frac{\mu_G(H) |\mathrm{Hom}(\Gamma, H)|}{|N_G(H)|}. \tag{5}$$

For example, it follows that for the Suzuki groups G we have

$$d_2(G) = \frac{1}{e} \sum_{f|e} \mu\left(\frac{e}{f}\right) 2^f (2^{4f} - 2^{3f} - 9),$$

where μ is the classical Möbius function on $\mathbb{N}$, given by $\mu(n) = (-1)^k$ if n is a product of k distinct primes, and $\mu(n) = 0$ otherwise. See [5] for this result, and for a number of other applications of Hall's theory to these groups G, and see Section 6 for an application to certain Hecke groups Γ.

This paper, together with its applications in [5], extends work by Silver and the second author [13], where orientably regular maps of type $\{4, 5\}$ with automorphism group $G \cong Sz(q)$ were enumerated, and by Hubard and Leemans [10], where regular maps and polytopes were enumerated. In each case the authors used a restricted form of Möbius inversion, concentrating mainly on subgroups $H \cong Sz(2^f)$ where f divides e. Here we compute $\mu_G(H)$ for all subgroups $H \leq G$, allowing the enumeration of a wider range of regular objects, including dessins d'enfants, regular or orientably regular maps and hypermaps, and coverings of topological spaces (see [5]).

The method used for calculating the values of μ_G is as follows. Hall [8, Theorem 2.3] showed that, in any finite group G, a subgroup H satisfies $\mu_G(H) = 0$ unless H is an intersection of maximal subgroups of G. In our case, with $G = Sz(q)$, instead of directly determining the set $\mathcal{I}$ of such intersections, we first describe, in Section 2.3.3, a more convenient set $\mathcal{S}$ of subgroups of G such that every subgroup in $\mathcal{I}$ is conjugate to a subgroup in $\mathcal{S}$ (see Theorem 3.1). Since μ_G is invariant under conjugation, it sufficient to find its values on $\mathcal{S}$; then the set $\mathcal{T}$ appearing in equation (5) is simply the subset of $\mathcal{S}$ on which μ_G can take non-zero values.

For each pair of subgroups $H, K \in \mathcal{S}$, we determine in Table 2 (Section 5) the number $N(H; K)$ of conjugates of K containing H. Since $\mu_G(K) = 0$ for all $K \notin \mathcal{S}$, equation (2) gives

$$\mu_G(H) = -\sum_{H<K\in\mathcal{S}} N(H; K)\mu_G(K) \tag{6}$$

where $H \in \mathcal{S} \setminus \{G\}$. This allows $\mu_G(H)$ to be calculated recursively, starting with $\mu_G(G) = 1$ and then using the values of $\mu_G(K)$ for the subgroups $K \in \mathcal{S}$ properly containing each $H \in \mathcal{S} \setminus \{G\}$.

Acknowledgements The authors are grateful to Dimitri Leemans for some very helpful comments on enumeration with Suzuki groups, and to Nikos Kanakis for help in preparing the TeX file.

2 The Suzuki groups and their subgroups

This section is based mainly on Suzuki's description of the groups $Sz(q)$ in [15]; see also [11, §XI.3] and [16, §4.2] for further information. We have largely followed Suzuki's notation for elements and subgroups, except that we use the symbol F for the subgroup denoted in [15] by H (a Frobenius group of order $q^2(q-1)$), while we use H for an arbitrary subgroup of G. Also, our rule for distinguishing the subgroups A_1 and A_2 in §2.1 is non-standard.

2.1 The definition of the Suzuki groups

Let $\mathbb{F} = \mathbb{F}(e)$ be the finite field $\mathbb{F}_q$ of $q = 2^e$ elements for some odd $e \geq 1$, and let θ be the automorphism $\alpha \mapsto \alpha^r$ of $\mathbb{F}$ where $r = \sqrt{2q} = 2^{(e+1)/2}$, so that θ^2 is the Frobenius automorphism $\alpha \mapsto \alpha^2$.

For any $\alpha, \beta \in \mathbb{F}$ let (α, β) denote the 4×4 matrix

$$(\alpha,\beta) = \begin{pmatrix} 1 & & & \\ \alpha & 1 & & \\ \alpha^{\theta+1}+\beta & \alpha^\theta & 1 & \\ \alpha^{\theta+2}+\alpha\beta+\beta^\theta & \beta & \alpha & 1 \end{pmatrix}.$$

Since $(\alpha,\beta)(\gamma,\delta) = (\alpha+\gamma, \alpha\gamma^\theta+\beta+\delta)$, these matrices (α,β) form a group $Q = Q(e)$ of order q^2, with identity element $(0,0)$.

For each $\kappa \in \mathbb{F}^* := \mathbb{F} \setminus \{0\}$ let a_κ denote the 4×4 diagonal matrix with diagonal entries ζ_i where $\zeta_1^\theta = \kappa^{1+\theta}$, $\zeta_2^\theta = \kappa$, $\zeta_3 = \zeta_2^{-1}$ and $\zeta_4 = \zeta_1^{-1}$. These matrices form a cyclic group $A_0 = A_0(e) \cong \mathbb{F}^*$ of order $q-1$. Since

$$a_\kappa^{-1}(\alpha,\beta)a_\kappa = (\alpha\kappa, \beta\kappa^{1+\theta}),$$

the group $F = F(e)$ generated by Q and A_0 in $GL_4(q)$ is a semidirect product of a normal subgroup Q by a complement A_0, so it has order $q^2(q-1)$.

We define $G = G(e)$ to be the subgroup of $GL_4(q)$ generated by F and the 4×4 matrix with entries 1 on the minor diagonal and 0 elsewhere (denoted by τ in [15]). This is the Suzuki group associated with $\mathbb{F}_q$, usually denoted by $Sz(q)$ or ${}^2B_2(q)$. It is, in fact, the subgroup of the symplectic group $Sp_4(q) = B_2(q)$ fixed by a certain automorphism of order 2.

In its natural action $g : [v] \mapsto [vg]$ on the projective space $\mathbb{P}^3(\mathbb{F})$, G acts as a doubly transitive permutation group of degree q^2+1 on the ovoid

$$\Omega = \Omega(e) = \{[\alpha^{\theta+2}+\alpha\beta+\beta^\theta,\ \beta,\ \alpha,\ 1] \mid \alpha,\beta \in \mathbb{F}\} \cup \{\infty\} \subset \mathbb{P}^3(\mathbb{F}),$$

where

$$\infty := [1,0,0,0] \in \mathbb{P}^3(\mathbb{F}).$$

The subgroup G_∞ of G fixing ∞ is F. This acts as a Frobenius group on $\Omega \setminus \{\infty\}$: its Frobenius kernel is Q, acting regularly on $\Omega \setminus \{\infty\}$, and A_0 is a Frobenius complement $G_{\infty,\omega}$, fixing a second point

$$\omega := [0,0,0,1] \in \Omega$$

and acting semiregularly on $\Omega \setminus \{\infty, \omega\}$. Thus the stabiliser of any three points in Ω is the identity subgroup I, so G acts on Ω as a Zassenhaus group.

There are cyclic subgroups of G of mutually coprime odd orders

$$2^e \pm 2^{(e+1)/2} + 1 = q \pm r + 1,$$

contained in Singer subgroups of $GL_4(q)$: note that since $r = \sqrt{2q}$,

$$(q + r + 1)(q - r + 1) = q^2 + 1,$$

which divides $q^4 - 1$. Let us choose a pair of subgroups $A_1 = A_1(e), A_2 = A_2(e)$ of G of these two orders, indexed according to the rule

$$|A_1(e)| = a_1(e) := 2^e + \chi(e)2^{(e+1)/2} + 1,$$

$$|A_2(e)| = a_2(e) := 2^e - \chi(e)2^{(e+1)/2} + 1,$$

where $\chi(e) = 1$ or -1 as $e \equiv \pm 1$ or ± 3 mod (8). (This rule for indexing A_1 and A_2 differs from that used in [11, 15], where the rule is that $|A_1(e)| > |A_2(e)|$ for all e; the rule adopted here has an advantage which will be explained in §2.3.1.)

2.2 Basic properties of Suzuki groups

Here we record some basic properties of G; see [11, 15] for proofs.

1. G has order $q^2(q^2 + 1)(q - 1)$, and is simple if $e > 1$. (The group $G(1)$ is isomorphic to $AGL_1(5)$, of order 20.)

2. $\mathrm{Aut}\, G$ is a semidirect product of $\mathrm{Inn}\, G \cong G$ by a cyclic group of order e acting as the Galois group $\mathrm{Gal}\, \mathbb{F}$ on matrix entries, so $|\mathrm{Aut}\, G| = e|G|$.

3. Any two subgroups of G conjugate to Q intersect trivially, and any two subgroups conjugate to F have their intersection conjugate to A_0.

4. Q is a Sylow 2-subgroup of G of order q^2 and of exponent 4. The centre Z of Q consists of the identity and the involutions of Q (the matrices $(0, \beta)$ for $\beta \in \mathbb{F}$), with Z and Q/Z elementary abelian of order q.

5. $ZA_0 \cong F/Z \cong AGL_1(q)$, with A_0 acting regularly by conjugation on the non-identity elements of Z and of Q/Z.

6. The involutions of G are all conjugate, as are the cyclic subgroups of order 4; however an element of order 4 is not conjugate to its inverse.

7. All elements of G except those in a conjugate of Q have odd order. Each maximal cyclic subgroup of G of odd order is conjugate to A_0, A_1 or A_2; the intersection of any two of them is trivial.

8. A non-identity element of G has two fixed points on Ω, one fixed point, or none as it is conjugate to an element of A_0, of Q or of A_i for $i = 1, 2$, or, equivalently, as it has order dividing $q - 1$, 4 or $q^2 + 1$.

2.3 Some particular subgroups

Here we list some particular subgroups of G, in the anticipation that any subgroup H not conjugate to a member of the list satisfies $\mu_G(H) = 0$, and can therefore be ignored in the enumerations mentioned in Section 1. This list enables us to give a statement of all values of the Möbius function, prior to proving it.

2.3.1 Subgroups associated with subfields

If f divides e then restricting matrix entries to the subfield $\mathbb{F}(f)$ of $\mathbb{F}$ of order 2^f yields a subgroup $G(f) = Sz(2^f)$ of G. This acts doubly transitively, with degree $2^{2f} + 1$, on the subset $\Omega(f)$ of Ω defined over $\mathbb{F}(f)$. Since the point ∞ is defined over the prime field $\mathbb{F}(1)$, its stabiliser in $G(f)$ is $F(f) := F \cap G(f)$, which acts faithfully on $\Omega(f) \setminus \{\infty\}$ as a Frobenius group with kernel $Q(f) := Q \cap G(f) = \{(\alpha, \beta) \mid \alpha, \beta \in \mathbb{F}(f)\}$ and complement $A_0(f) := A_0 \cap G(f) = \{a_\kappa \mid \kappa \in \mathbb{F}(f)^*\}$. (Note that since r and $r + 1$ are coprime to $q - 1$, we have $a_\kappa \in G(f)$ if and only if $\kappa \in \mathbb{F}(f)^*$.) Let $Z(f)$ denote the centre of $Q(f)$, an elementary abelian group of order 2^f. If f and h are divisors of e, and f divides h, then

$$G(f) \leq G(h),\ F(f) \leq F(h),\ Q(f) \leq Q(h),\ Z(f) \leq Z(h),\ A_0(f) \leq A_0(h).$$

Indeed, each of these five families of subgroups forms a lattice isomorphic to the lattice $\Lambda(e)$ of divisors of e, a fact which is useful in evaluating μ_G.

Similarly, we would like to choose cyclic subgroups $A_i(f)$ of G of order

$$|A_i(f)| = a_i(f) = 2^f \pm \chi(f)2^{(f+1)/2} + 1$$

corresponding to A_i for each $i = 1, 2$ so that they also satisfy $A_i(f) \leq A_i(h)$ whenever f divides h. The following result allows this:

Lemma 2.1 *For each $i = 1, 2$ and each f dividing e there is a cyclic subgroup $A_i(f)$ of A_i of order $a_i(f)$.*

Proof. Since A_i is cyclic, it is sufficient to show that if f divides e then $a_i(f)$ divides $a_i(e)$ for $i = 1, 2$. Let $m := 2^{2f} + 1 = a_1(f)a_2(f)$. Then $2^j \equiv -2^{j-2f} \bmod (m)$ for all $j \geq 2f$, so

$$2^e \equiv -2^{e-2f} \equiv 2^{e-4f} \equiv \cdots \equiv (-1)^{(e-f)/2f} 2^f \bmod (m)$$

since e is odd. Now $(e - f)/2f$ is even or odd as $e/f \equiv \pm 1 \bmod (4)$, giving $2^e \equiv \pm 2^f \bmod (m)$ respectively. Similarly,

$$2^{(e+1)/2} \equiv (-1)^k\, 2^l \bmod (m)$$

where

$$k = \lfloor \frac{e+1}{4f} \rfloor \quad \text{and} \quad 0 \leq l = \frac{e+1}{2} - 2kf < 2f.$$

Now k is even if $e/f \equiv 1$ or 3 mod (8), and odd if $e/f \equiv -1$ or -3 mod (8), while $l = (f+1)/2$ or $(3f+1)/2$ as $e/f \equiv \pm 1$ mod (4).

In the case where $e/f \equiv 1$ mod (8), so that $\chi(e) = \chi(f)$, it follows that

$$a_i(e) = 2^e \pm \chi(e)2^{(e+1)/2} + 1 \equiv 2^f \pm \chi(f)2^{(f+1)/2} + 1 = a_i(f) \bmod (m)$$

for $i = 1, 2$; since $a_i(f)$ divides m it divides $a_i(e)$. The argument is essentially the same if $e/f \equiv -3$ mod (8), with a change of sign in the coefficient of $2^{(f+1)/2}$ balanced by a change of sign in $\chi(f)$. If $e/f \equiv 3$ or -1 mod (8) a similar argument, using the factorisation

$$-2^f \pm 2^{(3f+1)/2} + 1 = (1 \pm 2^{(f+1)/2})(2^f \mp 2^{(f+1)/2} + 1),$$

gives the result. □

In fact, since A_i is cyclic, this shows that if $f \mid h \mid e$ then $A_i(f) \leq A_i(h)$, so for each $i = 1, 2$ the subgroups $A_i(f)$, where f divides e, form a lattice isomorphic to $\Lambda(e)$. This explains our non-standard choice of indexing for these two sets of subgroups. Note, however, that $A_i(f)$ is now not necessarily a subgroup of $G(f)$, though it is conjugate to such a subgroup.

2.3.2 The normalisers of some subgroups

The normaliser B_0 of A_0 in G is a dihedral group of order $2(q-1)$; it is the subgroup $G_{\{\infty,\omega\}}$ of G preserving the subset $\{\infty, \omega\}$ of Ω, with its subgroup A_0 fixing these two elements and its involutions transposing them. Let us choose a particular involution $c \in B_0$ and, for each f dividing e, define

$$B_0(f) := \langle A_0(f), c \rangle \leq B_0,$$

a dihedral group of order $2(2^f - 1)$ (so $B_0(1) \cong C_2$). If $f > 1$, then $B_0(f)$ is self-normalising whereas the normaliser of $A_0(f)$ is B_0.

For $i = 1, 2$ the normaliser B_i of A_i in G is a semidirect product of A_i and a cyclic group of order 4 generated by an element c_i satisfying $c_i^{-1} a c_i = a^{2^e}$ for all $a \in A_i$. For each f dividing e let

$$B_i(f) := \langle A_i(f), c_i \rangle \leq B_i,$$

so $|B_i(f)| = 4a_i(f)$, with $B_2(1) \cong C_4$. If $i = 1$ or $f > 1$ then $B_i(f)$ is self-normalising, whereas the normaliser of $A_i(f)$ is B_i.

By their construction, these groups $B_i(f)$ are (abstract) Frobenius groups of degree $a_i(f)$, and they satisfy $B_i(f) \leq B_i(h)$ for $i = 0, 1$ and 2 whenever $f \mid h \mid e$.

2.3.3 An important set of subgroups

For each f dividing e, we have defined the following subgroups of G, with the symbols (f) usually omitted when $f = e$:

$$G(f),\ F(f),\ Q(f),\ Z(f),\ B_i(f),\ A_i(f) \quad (i = 0, 1, 2). \tag{7}$$

Let $\mathcal{S}$ denote the set consisting of the subgroups in (7) for all f dividing e. The conjugacy class in G of any of these groups will be denoted by changing the appropriate italic capital letter to the corresponding script capital; thus $\mathcal{G}(f), \mathcal{F}, \ldots$ denote the conjugacy classes containing $G(f)$, F, and so on.

We note the following coincidences, conjugacies (denoted by $\sim$) and isomorphisms:

$$G(1) \sim B_1(1) \cong AGL_1(5), \quad F(1) = Q(1) \sim B_2(1) \cong C_4,$$

$$B_0(1) \sim Z(1) \cong C_2, \quad A_2(1) = A_0(1) = I.$$

In addition, if 3 divides e then

$$B_1(1) = B_1(3), \quad A_1(1) = A_1(3) \cong C_5.$$

Apart from these, any two distinct terms in (7) represent non-conjugate subgroups of G. In view of their special role in the following calculations, we will denote the class $\mathcal{A}_2(1) = \mathcal{A}_0(1)$ by $\mathcal{C}_1$, the class $\mathcal{B}_0(1) = \mathcal{Z}(1)$ by $\mathcal{C}_2$, and the class $\mathcal{F}(1) = \mathcal{Q}(1) = \mathcal{B}_2(1)$ by $\mathcal{C}_4$, since these consist of the cyclic subgroups of G of orders $1, 2$ and 4.

2.4 The Möbius function of a Suzuki group

We can now present the main result of this paper in the form of Table 1, which gives the non-zero values of $\mu_G(H)$ for the subgroups H of G; any subgroups H not appearing in Table 1 (such as $Q(f)$ and $Z(f)$ for $f > 1$) satisfy $\mu_G(H) = 0$, and can therefore be ignored in applying equations such as (3). Because of the conjugacies listed in §2.3.3, some conjugacy classes appear 'under an alias': for instance $\mathcal{F}(1)$ appears as $\mathcal{B}_2(1)$, and if 3 divides e then $\mathcal{G}(1)$ and $\mathcal{B}_1(1)$ appear as $\mathcal{B}_1(3)$. In the second column, $a_i(f) = 2^f \pm \chi(f)2^{(f+1)/2} + 1$ for $i = 1, 2$ (see §2.1 and §2.3.1). In the third column, the values of $|N_G(H)|$ are given for applications of equation (5). In the final column, μ is the classical Möbius function on $\mathbb{N}$, defined by

$$\sum_{m|n} \mu(f) = \delta_{n,1}$$

for all $n \in \mathbb{N}$, with the consequence that $\mu(n) = (-1)^k$ or 0 as n is or is not a product of k distinct primes for some $k \geq 0$.

Our aim is to show that the final column of this table is correct, by proving the following theorem:

Theorem 2.2 *Let G be a Suzuki group $Sz(2^e)$ for some odd $e > 1$, and let H be a subgroup of G. If $\mu_G(H) \neq 0$ then $\mu_G(H)$ is as given by Table 1.*

Note that when H is the identity subgroup, the value of $\mu_G(H)$ is $|G|\mu(e)$, supporting a conjecture of Conder that this value is divisible by $|G|$ whenever G is an almost simple group, that is, $S \leq G \leq \mathrm{Aut}\, S$ for some non-abelian finite simple group S. By [2, 8] this is true when $G = L_2(q)$ or $PGL_2(q)$, and computer calculations have verified the conjecture when S is one of the smaller alternating or sporadic simple groups.

Conjugacy class of H	$\lvert H\rvert$	$\lvert N_G(H)\rvert$	$\mu_G(H)$
$\mathcal{G}(f),\ 1<f\mid e$	$2^{2f}(2^{2f}+1)(2^f-1)$	$\lvert H\rvert$	$\mu(e/f)$
$\mathcal{F}(f),\ 1<f\mid e$	$2^{2f}(2^f-1)$	$\lvert H\rvert$	$-\mu(e/f)$
$\mathcal{B}_0(f),\ 1<f\mid e$	$2(2^f-1)$	$\lvert H\rvert$	$-\mu(e/f)$
$\mathcal{A}_0(f),\ 1<f\mid e$	2^f-1	$2(q-1)$	$2\frac{(2^e-1)}{(2^f-1)}\mu(e/f)$
$\mathcal{B}_1(f),\ 1<f\mid e$	$4a_1(f)$	$\lvert H\rvert$	$-\mu(e/f)$
$\mathcal{B}_2(f),\ 1<f\mid e$	$4a_2(f)$	$\lvert H\rvert$	$-\mu(e/f)$
$\mathcal{B}_2(1)=\mathcal{C}_4$	4	$2q$	$-2^e\mu(e)$
$\mathcal{B}_0(1)=\mathcal{C}_2$	2	q^2	$-2^{2e-1}\mu(e)$
$\mathcal{A}_0(1)=\mathcal{C}_1$	1	$\lvert G\rvert$	$\lvert G\rvert\mu(e)$

Table 1: Information about the subgroups H with non-zero values for $\mu_G(H)$

3 Subgroups H with $\mu_G(H)\neq 0$

As a first step towards proving Theorem 2.2, in this section we find some necessary conditions for a subgroup H of G to satisfy $\mu_G(H)\neq 0$.

3.1 Maxint subgroups

If G is any finite group, we shall say that a subgroup H of G is *maxint* if it is the intersection of a set of maximal subgroups of G (when $H=G$ this set is empty). The set of maxint subgroups of G will be denoted by $\mathcal{I}$. Hall proved in [8, Theorem 2.3] that if $H\notin\mathcal{I}$ then $\mu_G(H)=0$, so in determining μ_G one may restrict attention to the subgroups $H\in\mathcal{I}$. Since μ_G is preserved under conjugacy, it is sufficient to consider a set of representatives of the conjugacy classes of subgroups in $\mathcal{I}$. The main step in the proof of Theorem 2.2 is to show that if G is a Suzuki group $G(e)$ then the set $\mathcal{S}$ defined in §2.3.3 contains such a set of representatives:

Theorem 3.1 *If $H\in\mathcal{I}$ then H is conjugate in G to a subgroup in $\mathcal{S}$, that is, $\mathcal{I}$ is contained in the union of the conjugacy classes*

$$\mathcal{G}(f),\ \mathcal{F}(f),\ \mathcal{Q}(f),\ \mathcal{Z}(f),\ \mathcal{B}_i(f),\ \mathcal{A}_i(f)$$

of subgroups of G, where f divides e and $i=0,1$ or 2.

The rest of this section is devoted to a proof of this theorem. We will use the following criterion for a subgroup H of G to be in $\mathcal{I}$. Let $\mathcal{M}$ denote the set of all maximal subgroups of G, and let $\mathcal{M}(H)$ denote the set of those containing a particular subgroup H of G. Then the following, valid for any finite group G, is evident:

Lemma 3.2 *Let $H \leq G$. Then*

$$H \leq \bigcap_{M \in \mathcal{M}(H)} M,$$

with equality if and only if $H \in \mathcal{I}$.

3.2 Maximal subgroups

We will systematically apply Lemma 3.2 to the various subgroups H of G, using the following result:

Proposition 3.3 *The set $\mathcal{M}$ of maximal subgroups of G is given by*

$$\mathcal{M} = \bigcup_{e/f \text{ prime}} \mathcal{G}(f) \cup \mathcal{F} \cup \mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2.$$

This result is an immediate consequence of the following classification, due to Suzuki [15, Theorems 9 and 10]:

Proposition 3.4 *If $H \leq G$ then either $H \in \mathcal{G}(f)$ for some f dividing e, or H is a subgroup of a group in $\mathcal{F}$ or in $\mathcal{B}_i$ for some $i = 0, 1$ or 2.*

In the first case H is either simple or isomorphic to $G(1) \cong AGL_1(5)$, and in the second case H is solvable. Finite solvable groups H all satisfy Hall's theorems [7] on the existence and conjugacy of Hall π-subgroups for any set π of primes, generalising Sylow's theorems for single primes. We will use this fact, mainly with π the set $2'$ of odd primes.

Since $G(f) \in \mathcal{S}$ for each f dividing e, it follows from Proposition 3.4 that, in proving Theorem 3.1, it is sufficient to assume that H is a subgroup of a group in $\mathcal{F}$ or $\mathcal{B}_i$ for $i = 0, 1$ or 2. We will deal with these cases in turn.

In preparation for applying Lemma 3.2 in the first case, we will consider how the various maximal subgroups of G intersect F.

3.3 Point-stabilisers in maximal subgroups

Recall that F is the stabiliser in G of the point $\infty \in \Omega$. If $H \leq F$ then

$$\bigcap_{M \in \mathcal{M}(H)} M = \bigcap_{M \in \mathcal{M}(H)} (M \cap F),$$

so in applying Lemma 3.2 to H one can restrict attention to the stabilisers $M_\infty = M \cap F$ of ∞ for the various maximal subgroups M of G. The following result describes the possibilities for these stabilisers.

Lemma 3.5 *Let M be a maximal subgroup of G.*

1. *If $M = F^g \in \mathcal{F}$, then $M \cap F = F$ or $M \cap F = G_{\infty,\infty g} \in \mathcal{A}_0$ as $g \in F$ or not.*

2. *If $M = G(f)^g \in \mathcal{G}(f)$ for some $f|e$, then $M \cap F = F(f)^g \in \mathcal{F}(f)$ or $M \cap F = I \in \mathcal{C}_1$ as $g \in G(f)F$ or not.*

3. *If $M \in \mathcal{B}_0$ then $M \cap F \in \mathcal{A}_0 \cup \mathcal{C}_1 \cup \mathcal{C}_2$.*

4. *If $M \in \mathcal{B}_i$ for $i = 1, 2$ then $M \cap F \in \mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_4$.*

In order to prove part (2), we first need the following lemma:

Lemma 3.6 *If f divides e then $G(f)$ acts semi-regularly on $\Omega \setminus \Omega(f)$.*

Proof. By §2.2(8), a non-identity element of G fixes $2, 1$ or 0 elements of Ω as it has order dividing $q-1$, 4 or q^2+1. Similarly, a non-identity element of $G(f)$ fixes $2, 1$ or 0 elements of $\Omega(f)$ as it has order dividing $2^f - 1$, 4 or $2^{2f} + 1$ respectively. Since $2^f - 1$ divides $q - 1$, and $2^{2f} + 1$ divides $q^2 + 1$, a non-identity element of $G(f)$ can have no further fixed points in $\Omega \setminus \Omega(f)$. Thus all orbits of $G(f)$ on this set are regular, with point-stabilisers $G(f) \cap G_\alpha = I$ for $\alpha \in \Omega \setminus \Omega(f)$. □

Proof of Lemma 3.5. The maximal subgroups M of G are given by Proposition 3.3.

(1) This part is trivial, since F and M are the stabilisers in G of ∞ and ∞g, and G is doubly transitive on Ω.

(2) If $M = G(f)^g \in \mathcal{G}(f)$ then Lemma 3.6 shows that M acts doubly transitively on $\Omega(f)g$, and semiregularly on its complement. Thus $M \cap F = F(f)^g$ or I as $\infty \in \Omega(f)g$ or not, that is, as $g \in G(f)F$ or not.

(3) Each $M \in \mathcal{B}_0$ is the subgroup $G_{\{\alpha,\beta\}}$ of G preserving an unordered pair $\{\alpha, \beta\} \subset \Omega$. If $\infty \notin \{\alpha, \beta\}$ then since $G_{\alpha,\beta,\infty} = I$ we have $|M \cap F| \leq 2$, whereas if $\infty = \alpha$ or β then $M \cap F = G_{\alpha,\beta} \in \mathcal{A}_0$.

(4) If $M \in \mathcal{B}_i$ for $i = 1$ or 2 then $M = N_G(A) \cong A \rtimes C_4$ for some $A \in \mathcal{A}_i$; since A acts without fixed points, $M \cap F$ is isomorphic to a subgroup of C_4, so it is in $\mathcal{C}_m$ for $m = 1, 2$ or 4. □

3.4 Subgroups H of F

We first prove Theorem 3.1 for subgroups $H \in \mathcal{I}$ which are contained in groups in $\mathcal{F}$. Replacing H with a conjugate, we may assume that $H \leq F$.

3.4.1 Preliminaries

Here we record some observations and simplifications which will be used in the proof.

(a) Lemma 3.5 shows that each $M \in \mathcal{M}(H)$ satisfies $M \cap F \in \mathcal{X}_M$ where $\mathcal{X}_M = \mathcal{F}(f_M)$ for some f_M dividing e (depending on M), or $\mathcal{X}_M = \mathcal{A}_0$, or $\mathcal{X}_M = \mathcal{C}_m$ for some m dividing 4. If $\mathcal{X}_M = \mathcal{C}_m$ for some m, then since $H \leq M \cap F$ we have $H \in \mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_4$, so H is as required, i.e. conjugate to an element of $\mathcal{S}$; we may therefore assume that for each $M \in \mathcal{M}(H) \setminus \{F\}$ we have $\mathcal{X}_M = \mathcal{F}(f_M)$ for some f_M or $\mathcal{X}_M = \mathcal{A}_0$, with $M \in \mathcal{G}(f_M)$ or $\mathcal{F}$ respectively.

(b) Since F is solvable, Hall's theorems [7] imply that a Hall $2'$-subgroup A of H is contained in one of the Hall $2'$-subgroups of F. These are the conjugates of A_0, and Q permutes them regularly by conjugation, so by conjugating H with a suitable element of Q we may assume that $A \leq A_0$.

(c) If H has even order it contains an involution. The involutions in F (the non-identity elements of Z) are all conjugate under A_0, so in this case, by conjugating H with an element of A_0 we may also assume that H contains $z := (0, 1)$.

(d) If any $M \in \mathcal{M}(H) \setminus \{F\}$ satisfies $M = G(f_M)^g \in \mathcal{G}(f_M)$ for some $g \in G$, then since $M \cap F \in \mathcal{F}(f_M)$ we have $g \in G(f_M)F$ by Lemma 3.5. Without loss of generality we can therefore choose this conjugating element g to be in F. Then

$$F(f_M)^g = (G(f_M) \cap F)^g = G(f_M)^g \cap F = M \cap F.$$

Thus $z \in M \cap F$, so $z^{g^{-1}} \in F(f_M)$. Since $F = A_0Q$ we can write $g = ab$ where $a \in A_0$ and $b \in Q$. Since z is in the centre Z of Q we have

$$z^{g^{-1}} = (z^{b^{-1}})^{a^{-1}} = z^{a^{-1}},$$

so $z^{a^{-1}} \in F(f_M)$. Since $z \in F(1)$ and A_0 acts regularly by conjugation on the involutions in Z, this implies that $a \in A_0(f_M)$. Thus $g = ab$ with $a \in F(f_M)$, so each $M \in \mathcal{M}(H) \setminus \{F\}$ satisfies

$$M \cap F = F(f_M)^g = F(f_M)^b$$

for some f_M dividing e, with $b \in Q$.

(e) We claim that if $f \mid h \mid e$ then the set

$$Q(f, h) := \{g \in Q \mid Q(f)^g \leq Q(h)\}$$

is the union of the cosets $(\alpha, 0)Z$ of Z in Q where $\alpha \in \mathbb{F}(h)$.

Clearly this set consists of complete cosets of Z in Q. The elements $(\alpha, 0)$ where $\alpha \in \mathbb{F}$ are representatives of these cosets, since there is an epimorphism $(\alpha, \beta) \mapsto \alpha$ from Q to the additive group of $\mathbb{F}$, with kernel Z. Therefore it suffices to show that $g := (\alpha, 0) \in Q(f, h)$ if and only if $\alpha \in \mathbb{F}(h)$.

If $\alpha \in \mathbb{F}(h)$ then $g \in Q(h)$; since $Q(f) \leq Q(h)$ we have $Q(f)^g \leq Q(h)$ and hence $g \in Q(f, h)$. For the converse, note that $(1, 0) \in Q(f)$. A simple calculation shows that

$$(1, 0)^g = (1, \alpha + \alpha^\theta),$$

so if $g \in Q(f, h)$ then $\alpha + \alpha^\theta \in \mathbb{F}(h)$. The function $\phi : x \mapsto x + x^\theta$ maps each subfield $\mathbb{K}$ of $\mathbb{F}$ into itself. Composing ϕ with itself gives a quadratic polynomial

$$\phi^2 : x \mapsto (x + x^\theta) + (x + x^\theta)^\theta = x + x^{\theta^2} = x + x^2$$

defined over the prime field, so if $\beta \in \mathbb{K}$ then any element of $\phi^{-2}(\beta)$ has degree at most 2 over $\mathbb{K}$. Since e is odd, $\mathbb{F}$ contains no quadratic extensions, so $\phi^{-2}(\mathbb{K}) \subseteq \mathbb{K}$ and hence $\phi^{-1}(\mathbb{K}) \subseteq \mathbb{K}$. In particular, since $\phi(\alpha) \in \mathbb{F}(h)$ we have $\alpha \in \mathbb{F}(h)$.

We can now start the case-by-case analysis of maxint subgroups $H \leq F$.

3.4.2 Subgroups $H \leq F$ which are not 2-groups

First suppose that H is not a 2-group, or equivalently the Hall $2'$-subgroup $A \leq A_0$ of H is not the identity subgroup, so that $C_G(A) = A_0$. For each $M \in \mathcal{M}$, A is contained in a maximal cyclic subgroup A_M of M, which has order $2^{f_M} - 1$ if $M \in \mathcal{G}(f)$, and order $2^e - 1$ if $M \in \mathcal{F}$. This subgroup A_M centralises A, so it is contained in A_0; thus $A = A_0(f_M)$, where we take $f_M = e$ when $M \in \mathcal{F}$, since this is the unique subgroup of A_0 of order $2^{f_M} - 1$. Now H is the intersection of these subgroups M, so A is the intersection of the corresponding subgroups A_M; it therefore has the form $A = A_0(f)$ where f is the highest common factor of the divisors f_M of e.

If H has odd order then this gives $H = A \in \mathcal{A}_0(f)$, one of the types allowed for in Theorem 3.1, so we may assume that H has even order. As noted in §3.4.1(c), this allows us to assume that $z \in H$. This also implies that each $M \in \mathcal{M}(H) \setminus \{F\}$ is in $\mathcal{G}(f_M)$ for some f_M, for otherwise $M \in \mathcal{F}$ and so H is a subgroup of a two-point stabiliser $M \cap F$, which has odd order. As shown in §3.4.1(d), it follows that such subgroups M satisfy $M \cap F = F(f_M)^b$ for some $b \in Q$.

We have $A_0(f_M) = A \leq A_M \leq M \cap F = F(f_M)^b$, so $A_0(f_M)$ and $A_0(f_M)^{b^{-1}}$ are both point-stabilisers in $F(f_M)$; because $F(f_M)$ acts as a Frobenius group on $\Omega(f)$, its kernel $Q(f_M)$ permutes these point-stabilisers regularly by conjugation, so $A_0(f_M) = A_0(f_M)^{b^{-1}c}$ for some $c \in Q(f_M)$. Thus the element $b^{-1}c$ of Q normalises $A_0(f_M)$, so it also normalises $C_G(A_0(f_M)) = A_0$. However, Q permutes the conjugates of A_0 regularly by conjugation (since it is also a Frobenius group), so $b^{-1}c = 1$ and hence $b = c \in Q(f_M)$.

Thus $M \cap F = F(f_M)^b = F(f_M)$ for each $M \in \mathcal{M}(H) \setminus \{F\}$, so H, being the intersection of such subgroups $F(f_M)$, together with F, has the form $F(f) \cap F = F(f)$ for some divisor f of e, giving $H \in \mathcal{F}(f)$ as required.

3.4.3 Subgroups $H \leq F$ which are 2-groups

Now suppose that H is a 2-group, so $H \leq Q$. By §3.4.1(a), for each $M \in \mathcal{M}(H) \setminus \{F\}$ either $M \cap F \in \mathcal{F}(f_M)$ for some f_M dividing e, or $M \cap F \in \mathcal{A}_0$, with $M \in \mathcal{G}(f_M)$ or $\mathcal{F}$ respectively. We may assume that $H \neq I$, so H has even order and hence (as in §3.4.2) the second possibility cannot arise. Thus $M \cap F = F(f_M)^b$ for some $b \in Q$, as shown in §3.4.1(d). As Q is normal in F, and is a Sylow 2-subgroup of F, we have $M \cap Q = Q(f_M)^b$; comparing centres, we see that $M \cap Z = Z(f_M)^b = Z(f_M)$ since $Z(f_M)$, being central in Q, is normalised by b. Thus if $H \leq Z$ then

$$H \;=\; \bigcap_{M \in \mathcal{M}(H)} M \;=\; \bigcap_{M \in \mathcal{M}(H)} (M \cap Z) \;=\; \bigcap_{M \in \mathcal{M}(H)} Z(f_M) \;=\; Z(f),$$

where f is the highest common factor of the integers f_M, so $H \in \mathcal{Z}(f)$.

We may therefore assume that $H \not\leq Z$, so H contains an element of order 4. Since F has a single conjugacy class of cyclic subgroups of order 4, we may assume that H contains the subgroup $Q(1) = \{1, z, y^{\pm 1}\}$ where $y := (1, 0)$. Then $Q(1) \leq H \leq M \cap Q = Q(f_M)^b$, so by §3.4.1(e) we have $b^{-1} \in (\alpha, 0)Z$ for some $\alpha \in \mathbb{F}(f_M)$. This shows that $M \cap Q =$

$Q(f_M)^b = Q(f_M)$ for each $M \in \mathcal{M}(H) \setminus \{F\}$, so taking the intersection over all such M gives $H = Q(f) \in \mathcal{Q}(f)$ where f is the highest common factor of the integers f_M.

3.5 Subgroups H of B_i

Now suppose that H is a subgroup of a group in $\mathcal{B}_i$ for some $i = 0, 1$ or 2, and is maxint. Without loss of generality we may assume that $H \leq B_i$.

3.5.1 Subgroups H of B_0

Suppose that $H \leq B_0$. The subgroup $A := H \cap A_0 = H \cap F$ is maxint, since H is, it is contained in F, and it has odd order, so by an argument in §3.4.2 we see that $A = A_0(f)$ for some f dividing e. Now H contains A with index at most 2, so either $H = A_0(f)$, or H is a dihedral subgroup of B_0 conjugate (since $|A_0|$ is odd) to $B_0(f)$. Thus H is in $\mathcal{A}_0(f)$ or $\mathcal{B}_0(f)$.

3.5.2 Subgroups H of B_i for $i = 1$ or 2

Suppose that $H \leq B_i$ where $i = 1$ or 2. Let $A := H \cap A_i$. If $|A| = 1$ then H is isomorphic to a subgroup of $B_i/A_i \cong C_4$, so H is in $\mathcal{C}_m$ for some $m = 1, 2$ or 4. We may therefore assume that $|A| > 1$. Any subgroup $M \in \mathcal{M}(H) \setminus \{B_i\}$ contains A, which has order dividing q^2+1, so it follows from the classification of the maximal subgroups in Proposition 3.3 that M must be in $\mathcal{B}_i$ or in $\mathcal{G}(f_M)$ for some f_M dividing e. The first possibility can be dismissed, since distinct subgroups in $\mathcal{B}_i$ have intersections of order dividing 4, so $M \in \mathcal{G}(f_M)$. We can now argue as in §3.4.2, by considering subgroups centralising A, to show that $A = A_i(f)$ for some f dividing e.

Now $|H : A|$ divides $|B_i : A_i| = 4$. If $|H : A| = 1$ then $H = A_i(f) \in \mathcal{A}_i(f)$, as required. If $|H : A| = 4$ then H is a subgroup of B_i of order $4a_i(f)$; all subgroups of this order are conjugate in B_i to $B_i(f)$, so $H \in \mathcal{B}_i(f)$. We will show that the remaining case $|H : A| = 2$, where $|H| = 2a_i(f)$, cannot arise.

In a Suzuki group $Sz(q)$, any subgroup K of order $2m$, where m divides $q^2 + 1$, is contained in a unique subgroup K^* of order $4m$. (This is because $A_i < KA_i < B_i$ for $i = 1$ or 2, up to conjugacy, and the complements for A_i in B_i have mutually trivial intersections.) Applying this to the subgroup $K = H$, firstly as a subgroup of G, and then as a subgroup of each of the Suzuki subgroups $M \cong G(f)$ in $\mathcal{M}(H)$, we see that there is a subgroup $H^* \leq B_i$, containing H with index 2, such that $H^* \leq M$ for all $M \in \mathcal{M}(H)$. Lemma 3.2 then shows that $H \notin \mathcal{I}$

This completes the proof of Theorem 3.1. □

4 Size of conjugacy classes

An important step in proving the statement of the Möbius function of G in Theorem 2.2 is to determine the number of conjugates of each subgroup $H \in \mathcal{S}$, equal to the index in

G of its normaliser $N_G(H)$. The orders of some of these normalisers are noted in Table 1.

Theorem 4.1 *Let f divide e. Then*

1. $N_G(G(f)) = G(f)$ *and* $|\mathcal{G}(f)| = |G|/|G(f)|$;
2. $N_G(F(f)) = F(f)$ *and* $|\mathcal{F}(f)| = |G|/|F(f)|$ *if* $f > 1$;
3. $|N_G(Q(f))| = 2^{e+f}(2^f - 1)$ *and* $|\mathcal{Q}(f)| = |G|/2^{e+f}(2^f - 1)$ *if* $f > 1$;
4. $N_G(Z(f)) = QA_0(f)$ *and* $|\mathcal{Z}(f)| = |G|/2^{2e}(2^f - 1)$ *if* $f > 1$;
5. $N_G(B_i(f)) = B_i(f)$ *and* $|\mathcal{B}_i(f)| = |G|/|B_i(f)|$ *if* $i = 1$, *or if* $i = 0$ *or* 2 *and* $f > 1$;
6. $N_G(A_i(f)) = B_i$ *and* $|\mathcal{A}_i(f)| = |G|/|B_i|$ *if* $i = 1$, *or if* $i = 0$ *or* 2 *and* $f > 1$;
7. $|N_G(B_2(1))| = 2q$ *and* $|\mathcal{B}_2(1)| = q(q^2 + 1)(q - 1)/2$;
8. $|N_G(B_0(1))| = q^2$ *and* $|\mathcal{B}_0(1)| = (q^2 + 1)(q - 1)$.

Proof. (1) Let $H = G(f)$ where f divides e. If $f > 1$, then since $N_G(G(f))$ contains $G(f)$ it cannot be solvable, so by Proposition 3.4 it must be conjugate to $G(h)$ for some multiple h of f. Since $G(h)$ is simple, we must have $h = f$ and $N_G(H) = H$, giving $|\mathcal{G}(f)| = |G|/|H|$. The case $f = 1$ is dealt with in (5), since $G(1)$ is conjugate to $B_1(1)$.

(4) It is convenient to prove (4) before (2) and (3). Let $f > 1$. Any element of G normalising $Z(f)$ must fix its unique fixed point ∞, so $N_G(Z(f)) \leq F$. By §2.2(4), $F = QA_0$. Now $Z(f)$ is centralised by Q since it lies in the centre Z of Q, and §2.2(5) implies that $N_G(Z(f)) \cap A_0 = A_0(f)$, so $N_G(Z(f)) = QA_0(f)$, of order $|Q|.|A_0(f)| = q^2(2^f - 1) = 2^{2e}(2^f - 1)$.

(3) Any element of G normalising $Q(f)$ must normalise its characteristic subgroup $Z(f)$, so $N_G(Q(f)) \leq N_G(Z(f)) = QA_0(f)$. Now $A_0(f) \leq F(f) \leq N_G(Q(f))$, and §3.4.1(e) shows that $N_Q(Q(f)) = \bigcup_\alpha (\alpha, 0)Z$ with the union over all $\alpha \in \mathbb{F}(f)$, so $N_G(Q(f))$ has order $|Z|.2^f.|A_0(f)| = 2^{e+f}(2^f - 1)$.

(2) Clearly $N_G(F(f)) \leq N_G(Q(f)) \leq QA_0(f)$, and $A_0(f) \leq N_G(F(f))$. Since Z acts semi-regularly on $\Omega \setminus \{\infty\}$, it acts semi-regularly by conjugation on the subgroups of F in the conjugacy class $\mathcal{A}_0$, so $N_G(F(f)) \cap Z = Z(f)$. Hence, using the proof of part (3), we see that $N_G(F(f)) \leq F \cap G(f) = F(f)$. Thus $F(f)$ is self-normalising.

(5, 6) See §2.3.2.

(7, 8) For $i = 0$ and 2 the subgroups in $\mathcal{B}_i(1)$ are cyclic groups of orders 2 and 4 respectively, so they are contained in Sylow 2-subgroups of G. There are $q^2 + 1$ Sylow 2-subgroups, each conjugate to Q and containing $q - 1$ subgroups of order 2, and containing $(q^2 - q)/2$ of order 4. Since distinct Sylow 2-subgroups have trivial intersection, there are $(q^2 + 1)(q - 1)$ and $(q^2 + 1)(q^2 - q)/2$ such subgroups in G. In each case such subgroups are all conjugate, so their normalisers have order q^2 and $2q$. □

5 Calculating values of μ_G

We can now complete the proof of Theorem 2.2 by calculating $\mu_G(H)$ for each subgroup $H \in \mathcal{S}$. In order to use equation (6) for this (see §1), we first need to know, for each pair of subgroups $H, K \in \mathcal{S}$, the number $N(H;K)$ of conjugates in G of K containing H. If $M(H;K)$ denotes the number of conjugates in G of H contained in K, then a simple double counting argument gives

$$M(H;K)M(K;G) = M(H;G)N(H;K) \tag{8}$$

for all $H, K \in \mathcal{S}$. This allows $N(H;K)$ to be determined from the values of the function M. Now $M(H;G) = |\mathcal{H}|$ and $M(K;G) = |\mathcal{K}|$, where $\mathcal{H}$ and $\mathcal{K}$ are the conjugacy classes of subgroups of G containing H and K, so these values are given by Theorem 4.1. The values of $M(H;K)$ for $K \neq G$ can be found by using arguments similar to those used in proving Theorem 4.1, so details are omitted.

	$G(h)$	$F(h)$	$Q(h)$	$Z(h)$	$B_0(h)$	$A_0(h)$
$G(f)$	1					
$F(f)$	1	1				
$Q(f)$	2^{e-h}	2^{e-h}	1			
$Z(f)$	$2^{2(e-h)}$	$2^{2(e-h)}$	2^{e-h}	1		
$B_0(f)$	1				1	
$A_0(f)$	$\frac{(2^e-1)}{(2^h-1)}$	$\frac{2(2^e-1)}{(2^h-1)}$			$\frac{(2^e-1)}{(2^h-1)}$	1
$B_1(f), f \geqslant 1$	1					
$B_2(f)$	1					
$A_1(f), f \geqslant 1$	$\frac{a_1(e)}{a_1(h)}$					
$A_2(f)$	$\frac{a_2(e)}{a_2(h)}$					
$B_2(1) \cong C_4$	2^{e-h}	2^{e-h}	1			
$B_0(1) \cong C_2$	$2^{2(e-h)}$	$2^{2(e-h)}$	2^{e-h}	1	2^{2e-1}	

	$B_1(h)$	$B_2(h)$	$A_1(h)$	$A_2(h)$	$B_2(1)$	$B_0(1)$
$B_1(f), f \geqslant 1$	1					
$B_2(f)$		1				
$A_1(f), f \geqslant 1$	$\frac{a_1(e)}{a_1(h)}$		1			
$A_2(f)$		$\frac{a_2(e)}{a_2(h)}$		1		
$B_2(1) \cong C_4$	2^{e-1}	2^{e-1}			1	
$B_0(1) \cong C_2$	$2^{2(e-1)}$	$2^{2(e-1)}$			2^{e-1}	1

Table 2: Values of $N(H;K)$ where $H, K \in \mathcal{S}$.

The non-zero values of $N(H;K)$ resulting from (8) are given in Table 2, where the rows and columns are indexed by the subgroups H and K respectively; the row corresponding to the identity subgroup $H = A_0(1) = A_2(1)$ is omitted since in this case $N(H;K) = |\mathcal{K}|$, given by Theorem 4.1 for all $K \in \mathcal{S}$. The table is split into two parts, the second part giving further entries for the last six rows of the first part. We assume that f divides h and that $f > 1$ unless otherwise stated. Thus $G(1)$ is represented by its conjugate $B_1(1)$, while $F(1)$ and $Q(1)$ are represented by $B_2(1)$, and $Z(1)$ by $B_0(1)$ (see §2.3.3 and the comments in §2.4).

Given Table 2, one can systematically use equation (6) to calculate $\mu_G(H)$ for each $H \in \mathcal{S}$, starting with $H = G(f)$ in the first row, and working downwards through the table. For instance, if $H = G(f)$ then the subgroups $K \in \mathcal{S}$ with $N(H;K) \neq 0$ are those of the form $K = G(h)$ where $f \mid h \mid e$; under inclusion, these form a lattice isomorphic to the lattice $\Lambda(e/f)$ of all divisors h/f of e/f, with $\mu_G(K) = 1$ when $h = e$, so we find that $\mu_G(H) = \mu(e/f)$, as in Table 1. Next, if $H = F(f)$ we consider the subgroups $K = G(h)$ and $F(h)$ where $f \mid h \mid e$; these form a lattice isomorphic to $\Lambda(2e/f)$ since e is odd, giving $\mu_G(H) = \mu(2e/f) = -\mu(e/f)$. Similar arguments show that if $f > 1$ then $\mu_G(B_i(f)) = -\mu(e/f)$ for $i = 0, 1, 2$, and $\mu_G(Q(f)) = \mu_G(Z(f)) = \mu_G(A_i(f)) = 0$ for $i = 1, 2$. This process continues until $\mu_G(H)$ is evaluated for all $H \in \mathcal{S}$. The method is essentially the same as that described fully in [2, §4] for the groups $G = L_2(2^e)$, so the remaining details are omitted.

Since $\mu_G(H) = 0$ whenever $H = Q(f)$, $Z(f)$, $A_1(f)$ or $A_2(f)$ for any $f > 1$, we may disregard these subgroups, and let $\mathcal{T}$ denote the remaining set of subgroups $H \in \mathcal{S}$, namely those of the form

$$G(f),\ F(f),\ B_i(f)\ (i = 0, 1, 2),\ A_0(f),\ B_2(1),\ B_0(1),\ A_0(1), \tag{9}$$

where $1 < f \mid e$. This is a set of representatives for the conjugacy classes in Table 1; by the construction of $\mathcal{T}$, every subgroup H of G with $\mu_G(H) \neq 0$ must belong to one of these classes. This fact, together with the values of $|H|$, $|N_G(H)|$ and $\mu_G(H)$ determined earlier, justifies the entries in Table 1 and in particular proves Theorem 2.2. □

Each conjugacy class in Table 1 contains $|G|/|N_G(H)|$ subgroups, and $|\mathrm{Aut}\, G| = e|G|$ by §2.2(2), so equation (3) takes the form

$$n_\Gamma(G) = \frac{1}{e} \sum_{H \in \mathcal{T}} \frac{\mu_G(H)|\mathrm{Hom}(\Gamma, H)|}{|N_G(H)|},$$

as in equation (5). Table 1 gives the values of $\mu_G(H)$ and $|N_G(H)|$, so in order to apply this equation to a particular group Γ one needs only to count the homomorphisms $\Gamma \to H$ for each $H \in \mathcal{T}$. We will illustrate this in the case of certain Hecke groups Γ in Section 6.

6 Application to Hecke groups

In [2], the first author used Möbius inversion to enumerate the normal subgroups of the modular group $\Gamma = PSL_2(\mathbb{Z}) \cong C_2 * C_3$ with quotient group isomorphic to $L_2(q)$ for any

given prime power q. There are no normal subgroups of Γ with quotient group $G = Sz(q)$, since Suzuki groups have no elements of order 3, but instead one can apply the same method to Hecke groups other than Γ.

For each integer $k \geq 3$ the Hecke group H_k, introduced by Hecke in connection with Dirichlet series [9], is the subgroup of $PSL_2(\mathbb{R})$ generated by the Möbius transformations

$$X : z \mapsto z + \lambda_k \quad \text{and} \quad Y : z \mapsto \frac{-1}{z}$$

of the upper half plane, where $\lambda_k = 2\cos(\pi/k)$. This group is a free product of cyclic groups of orders 2 and k generated by Y and XY. In particular, H_3 is the modular group.

The torsion-free normal subgroups of H_k with a given finite quotient group G correspond bijectively to the orbits of $\operatorname{Aut} G$ on generating pairs for G of orders 2 and k. By equation (5) the number of these orbits is

$$\frac{1}{|\operatorname{Aut} G|} \sum_{H \leq G} \mu_G(H)|H|_2|H|_k \; = \frac{1}{e} \sum_{H \in \mathcal{T}} \frac{\mu_G(H)|H|_2|H|_k}{|N_G(H)|}, \tag{10}$$

where $|H|_n$ is the number of elements of order n in each subgroup H of G. Table 3 gives $|H|_n$ for subgroups $H \in \mathcal{T}$ in the cases $n = 2, 4$ and 5; here $q = 2^e$ and $1 < f \mid e$.

H	$\lvert H\rvert_2$	$\lvert H\rvert_4$	$\lvert H\rvert_5$
$G(f)$	$(2^f-1)(2^{2f}+1)$	$2^f(2^{2f}+1)(2^f-1)$	$2^{2f}(2^f-1)a_2(f)$
$F(f)$	2^f-1	$2^f(2^f-1)$	0
$B_0(f)$	2^f-1	0	0
$A_0(f)$	0	0	0
$B_1(f)$	$a_1(f)$	$2a_1(f)$	4
$B_2(f)$	$a_2(f)$	$2a_2(f)$	0
$B_2(1)$	1	2	0
$B_0(1)$	1	0	0
$A_0(1)$	0	0	0

Table 3: Values of $|H|_n$ for $n = 2, 4$ and 5.

With this information, together with Table 1, one finds from equation (10) that the numbers of (necessarily torsion-free) normal subgroups of H_4 and of H_5 with quotient group $Sz(q)$ are respectively

$$\frac{1}{e} \sum_{f|e} \mu\left(\frac{e}{f}\right) 2^f(2^f - 2) \tag{11}$$

and

$$\frac{1}{e} \sum_{f|e} \mu\left(\frac{e}{f}\right) (2^f - 1)a_2(f), \tag{12}$$

where $a_2(f) = |A_2(f)| = 2^f - \chi(f)2^{(f+1)/2} + 1$ (see §2.1).

One can apply similar arguments for odd $k > 5$. For instance, G contains elements of order 7 if and only if e is divisible by 3, in which case they form three conjugacy classes, represented by elements of A_0. It follows that the number of normal subgroups of H_7 with quotient group $Sz(q)$ is

$$\frac{3}{e}\sum_{3|f|e}\mu\left(\frac{e}{f}\right)(2^{2f}-2). \tag{13}$$

These enumerations have applications in other areas, such as topological graph theory. For instance, formulae (11), (12) and (13) give the numbers of orientably regular k-valent maps with automorphism group $Sz(q)$ for $k = 4, 5$ and 7 (see [5] for these and other examples).

References

[1] M. L. N. Downs, PhD thesis, University of Southampton, 1988.

[2] M. L. N. Downs, The Möbius function of $PSL_2(q)$, with application to the maximal normal subgroups of the modular group, *J. London Math. Soc.* 43 (1991), 61–75.

[3] M. L. N. Downs, The Möbius function of the Suzuki groups, preprint, 1993.

[4] M. L. N. Downs and G. A. Jones, Enumerating regular objects with a given automorphism group, *Discrete Math.* 64 (1987), 299–302.

[5] M. L. N. Downs and G. A. Jones, Enumerating regular objects associated with Suzuki groups, arXiv:1309.5215[math.GR], 2013.

[6] A. Grothendieck, Esquisse d'un Programme, in *Geometric Galois Actions I, Around Grothendieck's Esquisse d'un Programme* (ed. P. Lochak and L. Schneps) London Math. Soc. Lecture Note Ser. 242 (Cambridge University Press, Cambridge, 1997), 5–48.

[7] P. Hall, A note on soluble groups, *J. London Math. Soc.* 3 (1928), 98–105.

[8] P. Hall, The Eulerian functions of a group, *Q. J. Math.* 7 (1936), 134–151.

[9] E. Hecke, Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichung, *Math. Ann.* 112 (1936), 664–699.

[10] I. Hubard and D. Leemans, Chiral polytopes and Suzuki simple groups, in eds. R. Connelly, W. Whitely, A. Weiss, *Rigidity and Symmetry*, Fields Inst. Commun., to appear, 20 pages.

[11] B. Huppert and N. Blackburn, *Finite Groups III*, Springer-Verlag, Berlin – Heidelberg – New York, 1982.

[12] G. A. Jones, Enumeration of homomorphisms and surface-coverings, *Q. J. Math.* 46 (1995), 485–507.

[13] G. A. Jones and S. A. Silver, Suzuki groups and surfaces, *J. London Math. Soc.* (2) 48 (1993), 117–125.

[14] M. Suzuki, A new type of simple groups of finite order, *Proc. Nat. Acad. Sci. U.S.A.* 46 (1960), 868–870.

[15] M. Suzuki, On a class of doubly transitive groups, *Ann. of Math.* (2) 75 (1962), 105–145.

[16] R. A. Wilson, *The Finite Simple Groups*, Graduate Texts in Math. 251, Springer, London, 2009.